

Panserin 701

Complete Medium for Serum-free Cultivation of Lymphocytes from Whole Blood

Product	Description	Catalogue-No.	Size
Panserin 701	Ready-to-use serum-free medium for cultivation of lymphocytes	ST04-710701M ST04-710701	100 ml 500 ml

Product description

Panserin 701 is a complete, ready-to-use, serum-free medium for the cultivation of lymphocytes from whole blood.

Storage conditions

Storage: -20°C (in the dark)
Stability: 10 months from date of production
Size: 100 ml, 500 ml, other sizes on request

Composition

Based on Iscove's MEM the medium is enriched with additional trace elements, albumin, cholesterol, lipids and vitamins. Panserin 701 contains the mitogen phytohemagglutinin (PHA-L) for growth stimulation of lymphocytes.

Special advantages

Panserin 701 has been developed for the serum-free cultivation of lymphocytes from whole blood. The plant lectin (PHA) in Panserin 701 stimulates cell division.

Instructions for use

Isolation of lymphocytes from whole blood using density gradient centrifugation

- Mix heparinized blood 1:1 with DPBS (without Mg²⁺/Ca²⁺) and carefully transfer in centrifuge tubes, which were prefilled with lymphocyte separation medium (Pancoll, density 1.077 g/ml). Pipette carefully to avoid mixture of the different phases.
- Centrifuge the gradient at room temperature for 30 minutes at 800 g (brake of centrifuge switched "OFF"); centrifugation will reveal 4 phases:
 - Top yellow layer: plasma
 - Opaque white layer: lymphocytes
 - Clear layer: separation medium
 - Bottom layer: pellet with erythrocytes and granulocytes
- Remove the plasma with a pipette and transfer the lymphocytes with a fresh pipette into a new centrifuge tube.
- Wash the lymphocytes with DPBS (without Mg²⁺/Ca²⁺) and centrifuge at 200 g for 10 minutes. Repeat washing procedure.

Cultivation and stimulation of lymphocytes

- Resuspend lymphocytes in Panserin 701 at a cell density of 1×10^5 .
- Incubate for 48 to 72 hours, depending on the nature and origin of lymphocytes and on future use.
- Re-stimulation with Panserin 701 after 1-2 weeks of culture may be necessary.

Technical support

For technical support, questions or remarks please contact your local PAN-Seratech partner or the technical department of PAN-Seratech via email (info@pan-seratech.com) or phone +49-8543-601630.

FOR RESEARCH USE ONLY! Not approved for human or animal diagnostic or therapeutic procedures.